Digitized by the Internet Archive
in 2026

https://archive.org/details/journal-of-chemical-thermodynamics_1993_25_contents

THE JOURNAL OF CHEMICAL THERMODYNAMICS

Volume 25, 1993

EXECUTIVE EDITOR

Professor M. L. McGlashan
Patoka, Fairwarp, Uckfield, East Sussex TN22 3DT, U.K.

EDITORS

Dr P. A. G. O'Hare
National Institute of Standards and Technology
Building 222
Gaithersburg, MD 20899
U.S.A.

Professor W. A. Wakeham
Dept of Chem. Engineering and Chem. Technology
Imperial College
London SW7 2BY
U.K.

Professor R. D. Weir
Dept of Chemistry and Chem. Engineering
Royal Military College
Kingston
Ontario, Canada K7K 5L0

ADVISORY BOARD

Chairman: G. Somsen (The Netherlands)

E. H. P. Cordfunke (The Netherlands)
G. Della Gatta (Italy)
J. H. Dymond (U.K.)
M. B. Ewing (U.K.)
J. Fuger (Commission of the European Communities)
L. V. Gurvich (Russia)
Y. P. Handa (Canada)
H. F. Holmes (U.S.A.)
G. M. Olofsson (Sweden)
D. Patterson (Canada)
J. A. Rard (U.S.A)
G. M. Schneider (Germany)
M. Sorai (Japan)
W. V. Steele (U.S.A.)
F. L. Swinton (U.K.)
W. Wagner (Germany)

ACADEMIC PRESS
(Harcourt Brace & Company, Publishers)
London San Diego New York
Boston Sydney Tokyo Toronto

Copyright © 1993 Academic Press Limited

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 25

Number 1, January 1993

Goldberg; R. N.; Millero, F. J.; Tremaine, P. Loren G. Hepler and his scientific career 1

Pitzer, K. S. Thermodynamics of natural and industrial waters 7

Zhang, Y.; Wang, Y.; Shi, J.; Benson, G. C.; Lu, B. C.-Y. The enthalpy of dilution of calcium chloride in aqueous ethanol at the temperatures 288.15 K, 298.15 K, and 308.15 K 27

Jou, F.-Y.; Otto, F. D.; Mather, A. E. The solubility of propane in 1,2-ethanediol at elevated pressures 37

Grolier, J.-P. E.; Roux-Desgranges, G.; Berkane, M.; Jiménez, E.; Wilhelm, E. Heat capacities and densities of mixtures of very polar substances. 2. Mixtures containing *N*,*N*-dimethylformamide 41

Koga, Y.; Kristiansen, J.; Hvidt, Aa. Excess partial molar volumes of $\{xCH_3(CH_2)_3O(CH_2)_2OH + (1-x)H_2O\}$ in the water-rich region . . . 51

Garg, S. K.; Banipal, T. S.; Ahluwalia, J. C. Heat capacities and densities of liquid *o*-xylene, *m*-xylene, *p*-xylene, and ethylbenzene, at temperatures from 318.15 K to 373.15 K and at pressures up to 10 MPa 57

Apelblat, A. The vapour pressures of saturated aqueous lithium chloride, sodium bromide, sodium nitrate, ammonium nitrate, and ammonium chloride at temperatures from 283 K to 313 K 63

Larson, J. W.; Tewari, Y. B.; Goldberg, R. N. Thermochemistry of the reactions between adenosine, adenosine 5′-monophosphate, inosine, and inosine 5′-monophosphate; the conversion of L-histidine to (urocanic acid + ammonia) 73

Xu, Y.; Hepler, L. G. Calorimetric investigations of crystalline, molten, and supercooled $Ca(NO_3)_2 \cdot 4H_2O$ and of concentrated $Ca(NO_3)_2$(aq) 91

Holmes, H. F.; Mesmer, R. E. Isopiestic studies of $NaHSO_4$(aq) at elevated temperatures. Thermodynamic properties 99

Hopkins, H. P.; Hamilton, D. D.; Wilson, W. D.; Campbell, J.; Fumero, J. Effects of C_2H_5OH, Na^+(aq), $N(CH_2CH_3)_4^+$(aq), and Mg^{2+}(aq) on the thermodynamics of double-helix-to-random-coil transitions of poly(dA)*poly(dT) and poly(dAdT) . 111

Crovetto, R.; Lvov, S. N.; Wood, R. H. Vapor pressures and densities of NaCl(aq) and KCl(aq) at the temperature 623 K and $CaCl_2$(aq) at the temperatures 623 K and 643 K 127

Pagé, M.; Huot, J.-Y.; Jolicoeur, C. A thermodynamic investigation of aqueous 2-methoxyethanol 139

Roy, R. N.; Roy, L. N.; Davis, W. B.; Larkin, D. W.; Willard, D.; Johnson, D. A.; Millero, F. J. The thermodynamic behavior of hydrogen bromide in aqueous solutions of barium bromide 155

Hovey, J. K.; Pitzer, K. S.; Rard, J. A. Thermodynamics of Na_2SO_4(aq) at temperatures T from 273 K to 373 K and of $\{(1-y)H_2SO_4 + yNa_2SO_4\}$(aq) at $T = 298.15$ K . 173

Number 2, February 1993

Yoon, J.-H.; Lee, H.-S.; Lee, H. Solubilities of 2,4-dichloro-1-naphthol in supercritical carbon dioxide 193

Antosik, M.; Stryjek, R. (Liquid + liquid) equilibria in (hexadecane + tetralin + 3-methoxypropionitrile) 197

Matilla, A. D.; Tardajos, G.; Aicart, E. Thermodynamic mixing properties of (chlorobenzene + an alkane) 201

Callanan, J. E.; Weir, R. D.; Westrum, E. F., Jr. Thermodynamics of divalent-metal fluorides. VI. Heat capacity and thermodynamic properties of the fast-ion conductor $CaSn_2F_6$ at temperatures from 6 K to 310 K 209

Hatem, G.; Gaune-Escard, M. Calorimetric investigation of $\{xKF+(1-x)NdF_3\}$(l) . 219

Ribeiro da Silva, M. A. V.; Ferrão, M. L. C. C. H.; Lopes, A. J. M. Enthalpies of combustion of each of the two bromonaphthalenes 229

Marceca, E.; Fernández-Prini, R. Solubility of solids in near-critical fluids. II. CHI_3 and CI_4 in ethene 237

Abdulagatov, I. M.; Bazaev, A. R.; Ramazanova, A. E. Volumetric properties and virial coefficients of (water + methane) 249

Carson, A. S.; Laye, P. G.; Pedley, J. B.; Welsby, A. M. The enthalpies of formation of iodomethane, diiodomethane, triiodomethane, and tetraiodomethane by rotating combustion calorimetry 261

Konings, R. J. M.; Cordfunke, E. H. P.; Ouweltjes, W. The standard molar enthalpies of formation of CdI_2(s) and Cs_2CdI_4(s) 271

Tillner-Roth, R.; Baehr, H. D. Measurements of liquid, near-critical, and supercritical (p, ρ, T) of 1,1,1,2-tetrafluoroethane (R 134a) and of 1,1-difluoroethane (R 152a) . 277

Tewari, Y. B.; Kishore, N.; Margolis, S. A.; Goldberg, R. N.; Shibatani, T. Thermochemistry of the hydrolysis of L-arginine to (L-citrulline + ammonia) and of the hydrolysis of L-arginine to (L-ornithine + urea) 293

Ott, J. B.; Sipowska, J. T.; Gruszkiewicz, M. S.; Woolley, A. T. Excess volumes for (ethanol + water) at the temperatures (298.15 and 348.15) K and pressures (0.4, 5, and 15) MPa and at the temperature 323.15 K and pressures (5 and 15) MPa . 307

Advice to authors 319

Number 3, March 1993

López, M.; Legido, J. L.; Jiménez, E.; Romaní, L.; Carballo, E.; Paz Andrade, M. I. Excess molar enthalpies for $\{x_1CH_3(CH_2)_2OH + x_2CH_3CH_2COCH_2CH_3 + (1-x_1-x_2)CH_3(CH_2)_uCH_3\}$ ($u = 5, 6$) at the temperature 298.15 K 321

Miyanaga, S.; Chubachi, K.; Nakamura, M.; Tamura, K.; Murakami, S. Excess molar enthalpies and excess molar heat capacities of $[xCF_3CH_2OH + (1-x)\{HCON(CH_3)_2$ or $CH_3CN\}]$ at the temperature 298.15 K 331

Pintos, M.; Amigo, A.; Bravo, R. Effect of alkane chain-length on the excess volume of a binary mixture containing a cyclic ether 337

Cordfunke, E. H. P.; Konings, R. J. M.; Laan, R. R. van der; Ouweltjes, W. The thermochemical properties of $Ba_3B_2O_6$(s) 343

Hedwig, G. R.; Høiland, H. Thermodynamic properties of peptide solutions. 9. Partial molar isentropic pressure coefficients in aqueous solutions of sequence isomeric tripeptides with a single $-CH_3$ side-chain 349

Czarnota, I. Heat capacity of octane at high pressures 355

Malhotra, R.; Price, W. E.; Woolf, L. A. Thermodynamic properties of pentan-3-one at temperatures from 278 K to 338 K and pressures from 0.1 MPa to 380 MPa . 361

Góralski, P. A calorimetric study of cholesterol dissolved in an alcohol—the solute—solvent interaction 367

Pardo, J.; Rodríguez, V.; Lafuente, C.; Royo, F. M.; Urieta, J. S. Excess molar volumes of (an alkanediol + an alkanol) and of (dichloromethane + an alkanol or an alkanediol) at the temperature 303.15 K 373

Letcher, T. M.; Nevines, J. A.; Vijayan, R. P.; Radloff, S. E. Excess enthalpies of (a chlorinated benzene + an alkanol) at the temperature 298.15 K 379

Gruszkiewicz, M. S.; Ott, J. B.; Sipowska, J. T. Excess enthalpies for (propane + chloromethane) at the temperatures (298.15, 323.15, and 348.15) K and pressures (5 and 15) MPa, and at the temperature 363.15 K and pressures (5, 10, and 15) MPa 385

O'Hare, P. A. G. Calorimetric measurements of the specific energies of reaction of arsenic and of selenium with fluorine. Standard molar enthalpies of formation $\Delta_f H^\circ_m$ at the temperature 298.15 K of AsF_5, SeF_6, As_2Se_3, As_4S_4, and As_2S_3. Thermodynamic properties of AsF_5 and SeF_6 in the ideal-gas state. Critical assessment of $\Delta_f H^\circ_m(AsF_3, l)$ and the dissociation enthalpies of As–F bonds . . 391

Jackowski, A. W. The effect of polymorphic forms of tricosane on the phase equilibria of (pentane + tricosane) 403

Lukas, W. (Solid + liquid) equilibria of {potassium rhenate(VII) + rubidium rhenate(VII)} and {potassium rhenate(VII) + caesium rhenate(VII)} 409

Morss, L. R.; Day, P. P.; Felinto, C.; Brito, H. Standard molar enthalpies of formation of Y_2O_3, Ho_2O_3, and Er_2O_3 at the temperature 298.15 K 415

Ewing, M. B.; Goodwin, A. R. H. Speeds of sound, perfect-gas heat capacity, and second acoustic virial coefficient for air at the temperature 255 K and pressures in the range 0.031 MPa to 6.9 MPa 423

Shukla, N. K.; Prasad, R.; Sood, D. D. The standard molar enthalpies of formation at the temperature $T = 298.15$ K of barium molybdate $BaMoO_4$(cr) and strontium molybdate $SrMoO_4$(cr) 429

Finch, A.; Gardner, P. J.; Head, A. J.; Wu Xiaoping. The standard enthalpies of formation of the aqueous tetramethylammonium ion and tetraethylammonium ion 435

Krombach, R.; Schneider, G. M. Investigations of the high-pressure phase behaviour of thallium(I) tetradecanoate at pressures up to 250 MPa 445

Number 4, April 1993

Rowlinson, J. S. Thermodynamics of inhomogeneous systems. (Being the ninth ROSSINI LECTURE of the Commission on Thermodynamics of the International Union of Pure and Applied Chemistry delivered on Tuesday 18 August 1992 at its Conference on Chemical Thermodynamics held at Snowbird, UT, U.S.A., and printed here by permission of IUPAC.). 449

Suga, H. Frozen-in disorder and slow relaxation in crystals. (Being the 40th Huffman Memorial Award Lecture of the Calorimetry Conference delivered on 20 August 1992 at its 47th Conference organized jointly with the IUPAC Conference on Chemical Thermodynamics held at Snowbird, UT, U.S.A.) 463

Kabo, G. J.; Kozyro, A. A.; Krasulin, A. P.; Sevruk, V. M.; Ivashkevich, L. S. Thermodynamic properties and tautomerism of tetrazole 485

Ortega, J.; Gonzalez, E. Thermodynamic properties of (a methyl ester + an *n*-alkane). III. H_m^E and V_m^E for $\{xCH_3(CH_2)_{u-1}CO_2CH_3$ (u = 1 to 6) + $(1-x)CH_3(CH_2)_8CH_3\}$ 495

Tobaly, P.; Lanchec, G. Vapour pressure and enthalpy of sublimation of a series of organometallic complexes: $Cu(DPM)_2$, $Y(DPM)_3$, $Ba(DPM)_2$, and some derivatives 503

Ott, J. B.; Sipowska, J. T.; Woolley, A. T. Excess enthalpies for (propane + ethanol) at the temperatures (298.15, 323.15, 348.15, and 363.15) K and pressures (5, 10, and 15) MPa 511

Hallén, D. Enthalpies of solution and heat capacities for some α,ω-dichloroalkanes in water 519

Nakamura, M.; Chubachi, K.; Tamura, K.; Murakami, S. Excess molar volumes, excess isentropic and isothermal compressibilities, and excess molar isochoric heat capacities of $[xCF_3CH_2OH + (1-x)\{HCON(CH_3)_2$ or $CH_3CN\}]$ at the temperature 298.15 K 525

Viswanathan, R.; Darwin Albert Raj, D.; Lakshmi Narasimhan, T. S.; Balasubramanian, R.; Mathews, C. K. Study of (molybdenum + tellurium) by Knudsen-effusion mass spectrometry 533

Gee, N.; Freeman, G. R. Relative permittivities of 10 organic liquids as functions of temperature 549

García Baonza, V.; Fernández Rubio, J. E.; Cáceres Alonso, M.; Núñez Delgado, J. Thermodynamic properties of liquid carbon disulfide 555

Number 5, May 1993

González, E.; Ortega, J.; Matos, J. S.; Tardajos, G. Thermodynamic properties of (a methyl ester + an *n*-alkane). II. H_m^E and V_m^E for $\{xCH_3(CH_2)_{u-1}CO_2CH_3$ (u = 1 to 6) + $(1-x)CH_3(CH_2)_4CH_3\}$ 561

Iyoki, S.; Kuriyama, Y.; Tanaka, H.; Kira, K.; Okabe, T.; Uemura, T. Vapor-pressure measurements on (water + lithium chloride + lithium nitrate) at temperatures from 274.15 K to 463.15 K 569

Ribeiro da Silva, M. A. V.; Matos, M. A. R.; Monte, M. J. S.; Alves, M. C. B.; Vieira, J. M. A. P. Enthalpies of combustion, vapour pressures, and enthalpies of sublimation of 3-, 5-, 6-, and 8-aminoquinoline 579

Lebedev, B. V.; Kulagina, T. G. Thermodynamics of the $(CH_3)_2Zn \cdot S(CH_3)_2$ complex at temperatures from 0 to 330 K 591

Wormald, C. J.; Colling, C. N. Excess enthalpies of (water + argon)(g) at temperatures up to 698.2 K and pressures up to 13.3 MPa 599

Putnam, R. L.; Boerio-Goates, J. Heat-capacity measurements and thermodynamic functions of crystalline sucrose at temperatures from 5 K to 342 K. Revised values for $\Delta_f G^\circ_m$(sucrose, cr, 298.15 K), $\Delta_f G^\circ_m$(sucrose, aq, 298.15 K), S°_m(sucrose, aq, 298.15 K); and $\Delta_r G^\circ_m$(298.15 K) for the hydrolysis of aqueous sucrose 607

Fenclová, D.; Dohnal, V. Limiting activity coefficients for {1,1,2-trichlorotrifluoroethane (Freon-113) + an oxygen-containing substance or a hydrocarbon} . . . 615

Kiseleva, I. A.; Ogorodova, L. P.; Melchakova, L. V.; Bisengalieva, M. R. Thermodynamic properties of copper silicate: dioptase: $Cu_6Si_6O_{18} \cdot 6H_2O$ . . 621

Zakrzewski, M.; Handa, Y. P. Thermodynamic properties of ice and of tetrahydrofuran hydrate in confined geometries 631

Czarnota, I. Heat capacity of decane at high pressures 639

Salvi-Narkhede, M.; Adcock, J. L.; Gakh, A.; Van Hook, W. A. Vapor pressures, liquid molar volumes, vapor non-ideality, and critical properties of $CF_3OCF_2CF_2CF_3$, c-$CF_2CF_2CF_2CF_2O$, $CF_3OCF_2OCF_3$, and $CF_3OCF_2CF_2H$ 643

Kabata, Y.; Takada, M.; Sonobe, R.; Sugiura, I.; Uematsu, M. Properties of $\{xCF_3CH_2OH + (1-x)H_2O\}$ at temperatures from 310 K to 420 K. I. Bubble pressures and saturated-liquid densities 649

Acree, W. E., Jr.; Tucker, S. A.; Pilcher, G.; Paz Andrade, M. I.; Ribeiro da Silva, M. D. M. C. Enthalpies of combustion of *p*-azoxyanisole and *p*-azoxyphenetole: the dissociation enthalpy of the (N–O) bonds. Enthalpies of crystal-to-(liquid crystal) transitions 653

Treszczanowicz, T.; Cieślak, D. (Liquid + liquid) equilibria in (a dimethyl ether of a polyethene glycol + an *n*-alkane) 661

Ochel, H.; Becker, H.; Maag, K.; Schneider, G. M. Influence of a third component on (liquid + liquid) phase equilibria in (butan-2-ol + water) and in (butan-1-ol + water) at pressures up to 160 MPa 667

Rodríguez, V.; Lafuente, C.; López, M. C.; Royo, F. M.; Urieta, J. S. Excess molar volumes and vapour pressures of (benzene + each of several isomers of butanol). 679

BOOK REVIEW (Authors, Title, Reviewer)

Grenthe, I.; Fuger, J.; Konings, R. J. M.; Lemire, R. J.; Muller, A. B.; Cregu, C. N.-T.; Wanner, H. Chemical Thermodynamics of Uranium. **D. Cubicciotti** . . . 687

Number 6, June 1993

Fenclová, D.; Dohnal, V. (Vapour + liquid) equilibria of {1-bromo-1-chloro-2,2,2-trifluoroethane (halothane) + an oxygenated solvent or hexane} . . . 689

Kaji, K.; Tochigi, K.; Misawa, Y.; Suzuki, T. An adiabatic calorimeter for samples of mass less than 0.1 g and heat-capacity measurements on benzoic acid at temperatures from 19 K to 312 K 699

Tian, Y.-W.; Yu, H.-L.; Wang, Z.-C.; Zhang, X.-H.; Zhou, J.-K. High-temperature isopiestic studies on $\{(1-y)PbCl_2 + y(1-t)NaCl + ytBaCl_2\}$(l) at the temperature 873 K 711

Marceca, E.; Fernández-Prini, R. Solubility of solids in near-critical fluids. III. CHI_3 and CI_4 in sulphur hexafluoride 719

Chirico, R. D.; Knipmeyer, S. E.; Nguyen, A.; Smith, N. K.; Steele, W. V. The thermodynamic properties of 4,5,9,10-tetrahydropyrene and of 1,2,3,6,7,8-hexahydropyrene 729

Grønlund, F. Integrating the energy-balance equation in calorimetry: a way to simulate an adiabatic calorimeter 763

Mather, A. E.; Sadus, R. J.; Franck, E. U. Phase equilibria in (water + krypton) at pressures from 31 MPa to 273 MPa and temperatures from 610 K to 660 K and in (water + neon) from 45 MPa to 255 MPa and from 660 K to 700 K 771

Bayles, J. W.; Letcher, T. M.; Moollan, W. C. The determination of activity coefficients at infinite dilution using g.l.c. with moderately volatile solvents 781

McElroy, P. J.; Fang, Ji. Compression factors and second and third virial coefficients of (methane + ethene) using a direct method 787

Becker, P. J.; Schneider, G. M. (Liquid + liquid) and (liquid + liquid + gas) phase equilibria in (an alkan-1-ol + water) from (butan-1-ol + water) to (octadecan-1-ol + water) at pressures up to 200 MPa and temperatures up to 650 K 795

Number 7, July 1993

Gonzalez, E.; Ortega, J. Thermodynamic properties of (a methyl ester + an *n*-alkane). IV. H_m^E and V_m^E for $\{xCH_3(CH_2)_{u-1}CO_2CH_3$ $(u = 1$ to $6) + (1-x)CH_3(CH_2)_{10}CH_3\}$ 801

Róg, G.; Kozłowska-Róg, A.; Zakuła-Sokół, K.; Borchardt, G. Determination of the standard Gibbs free energies of formation of the calcium aluminates from the oxides by e.m.f. measurements 807

Barbarín-Castillo, J.-M.; McLure, I. A. Excess volumes of mixtures of nearly spherical molecules. 3. More mixtures containing octamethylcyclotetrasiloxane or decamethylcyclopentasiloxane 811

Abdulagatov, I. M.; Dvoryanchikov, V. I. Isochoric heat capacity of $\{xH_2O + (1-x)KOH\}$ near the critical point of pure water 823

Fenghour, A.; Wakeham, W. A.; Ferguson, D.; Scott, A. C.; Watson, J. T. R. Automated isochoric apparatus for the measurement of density of fluid mixtures at temperatures from 298.15 K to 773.15 K and pressures up to 40 MPa: results for helium and for nitrogen 831

Kishore, N.; Goldberg, R. N.; Tewari, Y. B. Apparent molar heat capacities and apparent molar volumes of aqueous glucose at temperatures from 298.15 K to 327.01 K 847

Liu, X.-Z.; Yang, J.-Z.; Ding, X.-H. Studies on the thermodynamics of the solvent extraction of metals. 4. {Diindium trisulfate + disodium sulfate + 2-ethylhexyl phosphoric acid mono-(2-ethylhexyl) ether + octane + water} 861

Nikitin, E. D.; Pavlov, P. A.; Skripov, P. V. Measurement of the critical properties of thermally unstable substances and mixtures by the pulse-heating method . . 869

Kabata, Y.; Takada, M.; Uematsu, M. Properties of $\{xCF_3CH_2OH + (1-x)H_2O\}$ at temperatures from 310 K to 420 K. II. Densities at pressures up to 200 MPa for $x = 0.8513$ 881

Rard, J. A. On the osmotic and activity coefficients of $Na_2WO_4(aq)$ and $Na_2MoO_4(aq)$ at the temperature 298.15 K, and the relations between mean activity coefficients of solutes under isopiestic conditions 887

Atik, Z.; Meents, B.; Gmehling, J. Excess molar enthalpies of (fluorobenzene or trifluoromethylbenzene or iodoethane + methanol or toluene) at the temperature 313.15 K and pressure 0.7 MPa 905

Zhu, S.; Shen, S.; Benson, G. C.; Lu, B. C.-Y. Excess enthalpies of (ethanol or propan-1-ol + cyclohexane + decane) at the temperature 298.15 K 909

O'Hare, P. A. G.; Hope, G. A.; Beck, C. M., II A new value for the standard molar enthalpy of formation $\Delta_f H_m^\circ$ of silicon sesquitelluride Si_2Te_3 at the temperature 298.15 K by combustion calorimetry in fluorine, and an assessment of the results of previous studies that lead to $\Delta_f H_m^\circ(Si_2Te_3)$ 919

Number 8, August 1993

Rudnyi, E. B.; Kaibicheva, E. A.; Sidorov, L. N. Negative ions VO_2^-, VO_3^-, $V_2O_5^-$, $V_3O_8^-$, and $V_4O_{10}^-$ in the vapors of vanadium oxides. Determining enthalpies of formation from equilibrium constants. 929

Wang Haijun; Zheng Guokang; Chen Mingzhi Excess molar volumes of (an aliphatic nitrile + an aromatic hydrocarbon) at the temperature 298.15 K. 949

Lebedev, B. V.; Kulagina, T. G.; Karatayev, Ye. N. Thermodynamics of the complex $(C_2H_5)_2Zn \cdot O(C_2H_5)_2$ at temperatures from 0 to 330 K 957

Steele, W. V.; Chirico, R. D.; Knipmeyer, S. E.; Nguyen, A. The thermodynamic properties of thianthrene and phenoxathiin. 965

Ramallo, A. V.; Legido, J. L.; Tojo, J. Densities, refractive indexes, and excess molar volumes of $\{x_1CH_3COCH_2CH_3 + x_2c\text{-}C_6H_{12} + (1-x_1-x_2)CH_3(CH_2)_5CH_3\}$ at the temperature 298.15 K 993

Sipowska, J. T.; Ott, J. B.; Woolley, A. T.; Marchant, B. G.; Gruszkiewicz, M. S. Excess enthalpies for (propane + methanol) at the pressure 5 MPa and the temperatures (253.15, 258.15, 263.15, and 273.15) K. The effect of water as an impurity on the (liquid + liquid) equilibria in (propane + methanol) 999

Kabata, Y.; Sonobe, R.; Sugiura, I.; Uematsu, M. Properties of $\{xCF_3CH_2OH + (1-x)H_2O\}$ at temperatures from 310 K to 420 K. III. Densities at pressures up to 200 MPa for $x = 0.6533$ 1005

Cordfunke, E. H. P.; Ouweltjes, W. The standard molar enthalpies of formation of Cs_2MnO_4, $Cs_2Si_4O_9$, and Cs_3PO_4 1011

Gruszkiewicz, M. S.; Ott, J. B.; Sipowska, J. T. Excess enthalpies for (ethane + chloromethane) at the temperatures (298.15, 323.15, 348.15, and 363.15) K and pressures from 5 MPa to 16 MPa 1017

Ren, H.; Yesavage, V. F. Enthalpy increments of $\{xCH_3OH + (1-x)(CH_3)_2CO\}$ between the temperatures 291 K and 610 K at pressures to 10 MPa 1031

Number 9, September 1993

Lorenzana, M. T.; Jiménez, E.; Legido, J. L.; Fernández, J.; Paz Andrade, M. I. Excess molar volumes of (1-chloropentane or 1-chlorohexane + heptane + decane) at the temperature 298.15 K. 1041

Singh, J.; Joshi, I. M.; Sharma, S. C. Excess molar volumes of (dibutyl ether + trichloromethane or tetrachloromethane or 1,2-dichloroethane or 1,1,2-trichloroethane or 1,1,2,2-tetrachloroethane or pentachloroethane) at the temperature 303.15 K 1049

Mussini, P. R.; Cipolli, A.; Mussini, T.; Rondinini, S. Standard aqueous potentials for the {thallium amalagam|thallium(I) iodide} electrode, and thermodynamic solubility product of thallium(I) iodide at temperatures from 298.15 K to 328.15 K . . 1055

Bruno, T. J.; Outcalt, S. L. Fugacity coefficients of hydrogen in (hydrogen + butane). 1061

Todorović, M.; Ninković, R. Osmotic and activity coefficients of $\{xKNO_3 + (1-x)K_2SO_4\}$(aq) at the temperature 298.15 K 1071

Zielkiewicz, J. (Vapour + liquid) equilibria in (heptane + ethanol + propan-1-ol) at the temperature 313.15 K 1077

Ortega, J.; Gonzalez, E. Thermodynamic properties of (a methyl ester + an *n*-alkane). V. H_m^E and V_m^E for $\{xCH_3(CH_2)_{u-1}CO_2CH_3$ ($u=1$ to 6) + $(1-x)CH_3(CH_2)_{12}CH_3\}$ 1083

Grønvold, F.; Stølen, S.; Tolmach, P.; Westrum, E. F., Jr. Heat capacities of the wüstites $Fe_{0.9379}O$ and $Fe_{0.9254}O$ at temperatures T from 5 K to 350 K. Thermodynamics of the reaction: $xFe(s) + (1/4)Fe_3O_4(s) = Fe_{0.7500+x}O(s) = Fe_{1-y}O(s)$ at $T \approx 850$ K, and properties of $Fe_{1-y}O(s)$ to $T = 1000$ K. Thermodynamics of formation of wüstite 1089

Miltenburg, J. C. van; Fourcade, J.; Ezzine, M.; Bergman, C. Thermodynamic properties of polymeric sulfur $S_{\omega 2}$ at temperatures between 5 K and 370 K 1119

López, E. R.; García, J.; Fernández, J.; Paz Andrade, M. I. Excess molar enthalpies of some examples of (a dichloroalkane + a ket-2-one) at the temperature 298.15 K . 1127

Grønvold, F. Enthalpy of fusion and temperature of fusion of indium, and redetermination of the enthalpy of fusion of tin 1133

Kabata, Y.; Takada, M.; Sonobe, R.; Sugiura, I.; Uematsu, M. Properties of $\{xCF_3CH_2OH + (1-x)H_2O\}$ at temperatures from 310 K to 420 K. IV. Densities at pressures up to 200 MPa for $x = 0.01935$, 0.05072, 0.1510, 0.3013, and 0.5011 . 1145

Fenghour, A.; Wakeham, W. A.; Ferguson, D.; Scott, A. C.; Watson, J. T. R. Densities of (water + nitrogen) in the temperature range 533 K to 703 K and pressures up to 31 MPa 1151

Korobov, M. V.; Savinova, L. N.; Sidorov, L. N. Stabilities of CoF_4 and CrF_5 in the gas phase. 1161

Number 10, October 1993

Diky, V. V.; Kabo, G. J.; Kozyro, A. A.; Krasulin, A. P.; Sevruk, V. M. Thermodynamic properties and pseudorotation of chlorocyclopentane 1169

Letcher, T. M.; Bricknell, B. C.; Sewry, J. D. Phase equilibria for (an alkanol + hept-1-yne + water) at the temperature 298.15 K 1183

Malhotra, R.; Woolf, L. A. Thermodynamic properties of 2-butoxyethanol at temperatures from 288 K to 348 K and pressures from 0.1 MPa to 380 MPa . 1189

Ngeyi, S. P.; Abhari, R.; Westrum, E. F., Jr. Thermodynamics of alkali alkanoates. IX. Heat capacity and thermodynamic functions of lithium 2-methylpropanoate at temperatures from 7 K to 535 K. 1197

Vis, M. G. M. van der; Cordfunke, E. H. P. The standard molar enthalpy of formation of tetraethoxysilane: $Si(OC_2H_5)_4(l)$ 1205

Huntelaar, M. E.; Cordfunke, E. H. P.; Ouweltjes, W. The standard molar enthalpies of formation of $SrZrSi_2O_7(s)$ and $Sr_6ZrSi_5O_{18}(s)$ 1211

Tufeu, R.; Subra, P.; Plateaux, C. Contribution to the experimental determination of the phase diagrams of some (carbon dioxide + a terpene) mixtures 1219

Hrubý, J.; Klomfar, J.; Šifner, O. (T, p, ρ) relation of liquid methanol at temperatures from 205 K to 321 K and pressures up to 50 MPa 1229

Zielkiewicz, J. Excess molar volumes of (heptane + ethanol + propan-1-ol) at the temperature 313.15 K 1243

Gospodinov, G. G.; Mihov, D. I. Molar heat capacities of the tellurites $MnTeO_3$, $MnTe_2O_5$, and $Mn_2Te_3O_8$ 1249

Acree, W. E., Jr., Tucker, S. A.; Pilcher, G.; Chowdhary, A.; Ribeiro da Silva, M. D. M. C.; Monte, M. J. S. Enthalpies of combustion of 2,2′,4,4′,6,6′-hexamethylazobenzene-*N*,*N*-dioxide, 2,2′,6,6′-tetramethylazobenzene-*N*,*N*-dioxide, 2,4,6-trimethylnitrobenzene, and 2,6-dimethylnitrobenzene: the dissociation enthalpies of the N=N and N–O bonds 1253

Ribeiro da Silva, M. A. V.; Monte, M. J. S.; Giera, E.; Kakolowicz, W. Standard molar enthalpy of formation of bis(2,2,6,6-tetramethylheptane-3,5-dionato) dioxouranium(VI) in the crystalline and gaseous states: the mean U–O bond-dissociation enthalpy 1263

Möller, D.; Gammon, B. E.; Marsh, K. N.; Hall, K. R.; Holste, J. C. Enthalpy-increment measurements from flow calorimetry of CO_2 and of $\{xCO_2 + (1-x)C_2H_6\}$ from pressures of 15 MPa to 18 MPa between the temperatures 230 K and 350 K . 1273

Number 11, November 1993

Nagata, I. Excess enthalpies of (aniline + butan-1-ol) and of (aniline + butan-1-ol + benzene) at the temperature 298.15 K 1281

Holstein, W. L. Thermodynamic properties of thallium(I) hydroxide in the vapor state. 1287

Araujo, M. E.; Maciel, M. R. W.; Francesconi, A. Z. Excess Gibbs free energies of (hexane + butan-2-ol) and of (cyclohexane + butan-2-ol) at the temperatures (323.15, 338.15, and 348.15) K 1295

Plesnar, Z; Bylicki, A. (Solid + liquid) equilibria in (hexadecan-1-ol + hexadecane) . 1301

Nakamura, M.; Chubachi, K.; Tamura, K.; Murakami, S. Thermodynamic properties of $[x\{HCON(CH_3)_2$ or $CH_3CN\} + (1-x)(CH_3)_2SO]$ at the temperature 298.15 K . 1311

Santos, L. C. R. dos; Oliveira, S. F. de; Espínola, J. G. P.; Airoldi, C. Thermochemistry of adducts of *N*,*N*-dimethylthioacetamide and of *N*,*N*-dimethylthioformamide with antimony triiodide or bismuth triiodide at the temperature 298.15 K . . 1319

Iglesias, T. P.; Legido, J. L.; Romaní, L.; Peleteiro, J.; Paz Andrade, M. I. Dielectric permittivities, densities, and excess molar volumes of $\{xCH_3(CH_2)_6CH_3 + (1-x)CH_3(CH_2)_vOH\}$ $(v = 2$ to $4)$ at the temperature 298.15 K 1325

O'Hare, P. A. G. Thermodynamic properties of silicides. III. Specific energy of combustion in fluorine of a hyperstoichiometric molybdenum disilicide. The standard molar enthalpy of formation $\Delta_f H_m^\circ$ of $MoSi_{2.067\pm0.002}$ at the temperature 298.15 K 1333

Yadav, B. L.; Maken, S.; Kalra, K. C.; Singh, K. C. Excess volumes of (an alkanol + an aromatic hydrocarbon) at the temperature 308.15 K 1345

Pavese, F. Recalculation on ITS-90 of accurate vapour-pressure equations for e-H_2, Ne, N_2, O_2, Ar, CH_4, and CO_2 1351

Ren, H.; Yesavage, V. F. Enthalpy increments of $\{xCH_3(CH_2)_3CH_3 + (1-x)(CH_3)_2CO\}$ between the temperatures 291 K and 610 K at pressures to 10 MPa . . . 1363

An Xueqin; Shen Weiguo; Wang Haijun; Zheng Guokang The (liquid + liquid) critical phenomena of (a polar liquid + an *n*-alkane). I. Coexistence curves of (*N*,*N*-dimethylacetamide + hexane) 1373

Finch, A.; Gardner, P. J.; Head, A. J.; Wu Xiaoping; Yefimov, M. E.; Furkaluk, M. U. The standard enthalpy of formation of the aqueous cyanide ion 1385

Callanan, J. E.; Weir, R. D.; Westrum, E. F., Jr. Thermodynamic properties of silicides. II. Heat capacity at temperatures *T* from 5.9 K to 341 K and derived thermodynamic properties to $T = 1200$ K of tungsten disilicide: $WSi_{2.06}$ . . 1391

Artigas, H.; Santafé, J.; López, M. C.; Royo, F. M.; Urieta, J. S. Excess volumes of (an alkanol + a haloalkane) at the temperatures 298.15 K and 313.15 K . . . 1403

Number 12, December 1993

Announcement i

Kozyro, A. A.; Kabo, G. J.; Krasulin, A. P.; Sevruk, V. M.; Simirsky, V. V.; Sheiman, M. S.; Frenkel, M. L. Thermodynamic properties of methylurea 1409

Tillner-Roth, R. An experimental study of the thermodynamic properties of the refrigerant mixture: {1,1,1,2-tetrafluoroethane (R 134a) + 1,1-difluoroethane (R 152a)} 1419

Apelblat, A. Solubilities of organic salts of magnesium, calcium, and iron in water . 1443

Plesnar, Z.; Bylicki, A. (Solid + liquid) equilibria in (dodecan-1-ol + dodecane) . 1447

Schmidt, J. A.; Sagua, A. E. Thermodynamic quantities for the ternary compound Stromeyerite: $Cu_{1+\delta}Ag_{1-\delta}S$ for $0 \leqslant \delta \leqslant 0.1$ 1453

Chirico, R. D.; Knipmeyer, S. E.; Nguyen, A.; Steele, W. V. The thermodynamic properties to the temperature 700 K of naphthalene and of 2,7-dimethylnaphthalene 1461

Ababio, B. D.; McElroy, P. J. (Pressure, amount-of-substance density, temperature) of $\{(1-x)CO_2 + xH_2\}$ using a direct method 1495

Ewing, M. B.; Goodwin, A. R. H. Speeds of sound in a natural gas of specified composition at the temperature 255 K and pressures in the range 64 kPa to 6.1 MPa 1503

Apelblat, A. The vapour pressures of saturated aqueous solutions of potassium bromide, ammonium sulfate, copper(II) sulfate, iron(II) sulfate, and manganese(II) dichloride, at temperatures from 283 K to 308 K 1513

NOTE

Barbarín-Castillo, J.-M.; McLure, I. A. The application of McGlashan and Potter's correlation for second virial coefficients to polymethyl substances . . . 1521

INDEX TO VOLUME 25 1523